



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## ALGEBRA.

113. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Find a square number consisting of 24 figures such that the numbers formed by the first 12 figures and the last 12 figures, respectively, are consecutive, and *vice versa*.

Solution by the PROPOSER.

(1). Let  $y^2 = 1000000000001x + 1000000000000$  be the number.

$$\therefore 1000000000001x = y^2 - 1000000000000.$$

$$\therefore x = \frac{(y+1000000)(y-1000000)}{10001 \times 99990001}.$$

$$\text{Let } y+1000000=99990001u, \quad y-1000000=10001v.$$

$$\therefore 10001v = 99990001u - 2000000. \quad v = 9998u - 199 + \frac{3u-9801}{10001}.$$

$$\text{Let } S = \frac{3u-9801}{10001}. \quad \therefore u = \frac{10001S+9801}{3} = 3333S + 3267 + \frac{2S}{3}.$$

$$\text{Let } S=3m. \quad \therefore u=10001m+3267.$$

$$y=1000000000001m+326666333267.$$

$$x=(99990001m+32663267)(10001m+3267).$$

$$\text{Let } m=0. \quad \therefore y=326666333267, \quad x=106710893289.$$

$$y^2=106710893290106710893289=1000000000000(x+1)+x.$$

$$(2). \quad y^2=1000000000001x+1, \text{ or } x=\frac{(y+1)(y-1)}{(10001)(99990001)}.$$

$$y+1=99990001u, \quad y-1=10001v.$$

$$\therefore v = \frac{9998u+(3u-2)}{10001} = 9998u + S. \quad \therefore u = \frac{3333S+(2S+2)}{3} = 3333S + t.$$

$$\therefore S=t-1+\frac{1}{3}t=t-1+m. \quad \therefore S=3m-1, \quad u=10001m-3333.$$

$$y=1000000000001m-333266673334.$$

$$x=(10001m-3333)(99990001m-33323335).$$

$$\text{Let } m=1.$$

$$\therefore y=666733326667, \quad x=444533328888.$$

$$y^2=444533328888444533328889=1000000000000x+(x+1).$$

114. Proposed by ELMER SCHUYLER, B. Sc., Professor of German and Mathematics in Boys' High School, Reading, Pa.

$$\left. \begin{aligned} bx^3 &= 10a^2bx + 3a^3y \\ ay^3 &= 10b^2ay + 3b^3x \end{aligned} \right\}$$

I. Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa., and the PROPOSER.

$$bx^3=10a^2bx+3a^3y\dots\dots(1), ay^3=10ab^2y+3b^3x\dots\dots(2).$$

Let  $x/a=u$ ,  $y/b=v$ , then (1) and (2) become

$$u^3=10u+3v\dots\dots(3), v^3=10v+u\dots\dots(4).$$

$$(3)+(4) \text{ gives } u^3+v^3-13(u+v)=0\dots\dots(5).$$

$$(3)-(4) \text{ gives } u^3-v^3-7(u-v)=0\dots\dots(6).$$

$$\text{From (5), } u+v=0\dots\dots(7); u^2-uv+v^2=13\dots\dots(8).$$

$$\text{From (6), } u-v=0\dots\dots(9), u^2+uv+v^2=7\dots\dots(10).$$

$$\text{From (7) and (9), } u=0, v=0.$$

$$(8)+(10) \text{ gives } u^2+v^2=10\dots\dots(11).$$

$$(8)-(10) \text{ gives } uv=-3\dots\dots(12).$$

$$\text{From (11) and (12), } u+v=\pm 2, u-v=\pm 4.$$

$$\therefore u=\pm 3 \text{ or } \pm 1, v=\mp 1 \text{ or } \mp 3.$$

$$\text{From (7) and (10), } u=\pm\sqrt{7}, v=\mp\sqrt{7}.$$

$$\text{From (9) and (8), } u=\pm\sqrt{13}, v=\pm\sqrt{13}.$$

$$\therefore u=0, 1, -1, 3, -3, \sqrt{7}, -\sqrt{7}, \sqrt{13}, -\sqrt{13}.$$

$$v=0, -3, 3, -1, 1, -\sqrt{7}, \sqrt{7}, \sqrt{13}, -\sqrt{13}.$$

$$x=0, a, -a, 3a, -3a, a\sqrt{7}, -a\sqrt{7}, a\sqrt{13}, -a\sqrt{13}.$$

$$y=0, -3b, 3b, -b, b, -b\sqrt{7}, b\sqrt{7}, b\sqrt{13}, -b\sqrt{13}.$$

II. Solution by COOPER D. SCHMITT, A. M., Professor of Mathematics, University of Tennessee, Knoxville, Tenn.

$$\text{Let } bx/ay=t.$$

$$\text{Divide (1) by (2), then } \frac{b^3x^3}{a^3y^3}=\frac{10bx+3ay}{10ay+3bx}, \text{ or } t^3=\frac{10t+3}{10+3t}.$$

$$\text{Whence, } 3t^4+10t^3-10t-3=0.$$

$$\text{Factoring, } (t^2-1)(3t^2+10t+3)=0.$$

$$\text{Whence, } t=1, -1, -\frac{1}{3}, -3.$$

$$\text{When } t=1, bx=ay \text{ and substituting in (1) we easily get,}$$

$$x=0, \text{ or } \pm a\sqrt{13}.$$

$$y=0, \text{ or } \pm b\sqrt{13}.$$

$$\text{When } t=-1, bx=-ay. \text{ and we get in a similar manner,}$$

$$x=0, \text{ or } \pm a\sqrt{7}.$$

$$y=0, \text{ or } \mp b\sqrt{7}.$$

$$\text{When } t=-\frac{1}{3},$$

$$x=0, \text{ or } \pm 3a.$$

$$y=0, \text{ or } \mp b.$$

$$\begin{aligned}\text{When } t &= -3, & x &= 0, \text{ or } \pm a. \\ & & y &= 0, \text{ or } \mp 3b.\end{aligned}$$

Also solved by *J. M. BOORMAN*, and *HARRY S. VANDIVER*.


## GEOMETRY.

143. Proposed by *J. T. FAIRCHILD*, A. M., Instructor in Mathematics, Crawfis College, Crawfis College, Ohio.

If the centers of three spheres do not lie in the same straight line, their surfaces cannot have more than two points in common. These points lie in a straight line perpendicular to the plane of centers and equal distances from this plane on opposite sides. [From *Phillips and Fisher's Geometry*.]

Solution by *G. B. M. ZERR*, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let  $A, B, C$  be the centers of the three spheres, respectively.  $B$  intersects  $A$  in the circle, radius  $FD$ , such that  $FD$  is perpendicular to  $AB$  and  $D$  lies on  $AB$ .  $C$  intersects  $A$  in the circle, radius  $EH$ , such that  $EH$  is perpendicular to  $AC$  and  $E$  lies on  $AC$ . The two circles  $D$  and  $E$  on the same sphere  $A$  can intersect in only two points situated on a line perpendicular to the plane  $AED$  and equally distant from it. But plane  $AED$  coincides with plane  $ABC$ . Therefore, the truth of the theorem follows.



144. Proposed by *L. C. WALKER*, Assistant in Mathematics, Leland Stanford, Jr. University, Palo Alto, Cal.

Find the equations of four cones that pass through three given straight lines intersecting in the same point.

I. Solution by the PROPOSER.

Let the mutual inclination of the line be  $2\alpha, 2\beta, 2\gamma$ , and let the equation of the cone be referred to the three given straight lines as coördinate axes.

The equation of the concentric sphere referred to the same axes is

$$x^2 + y^2 + z^2 + 2yz\cos 2\alpha + 2zx\cos 2\beta + 2xy\cos 2\gamma = r \dots (1).$$

The equation of the plane that passes through the intersection of the cone and (1) is

$$\frac{x}{r} + \frac{y}{r} + \frac{z}{r} = 1 \dots (2).$$

Now making (1) homogeneous by means of (2), and reducing, we have for the equation of the cone,